FS 2.60/1:12/32

Morbidity and Mortality Weekly Report

U. S. DEPARTMENT OF HEALTH, EDUCATION, AND WELFARE
PUBLIC HEALTH SERVICE

Prepared by the COMMUNICABLE DISEASE CENTER *634-5131*

For release August 16, 1963 — ATLANTA, GEORGIA 30333 — Vol. 12, No. 32

UNIVERSITY OF GEORGIA LIBRARIES AUG 1963

PROVISIONAL INFORMATION ON SELECTED NOTIFIABLE DISEASES IN THE UNITED STATES AND ON DEATHS IN SELECTED CITIES FOR WEEK ENDED AUGUST 10, 1963

POLIOMYELITIS – The 12 cases of poliomyelitis, 11 paralytic, reported for the week ending August 10, bring to a total of 153 the number of cases reported thus far in 1963. This figure remains well below one-half the level reported in 1962.

Five cases, all paralytic, were reported from Alabama. South Carolina and Illinois reported two cases each. Single cases occurred in five other States.

All of the Alabama cases have occurred in the northwest part of the State, with two cases in Lawrence County, and one each in Cullman, Madison, and Shelby Counties. Lawrence and Cullman Counties both border on Walker County, where 7 cases, with Type I poliovirus recovered from 2, have occurred (See MMWR, Vol. 12, pages 221, 229, 245, and 253).

Approximately 44,000 Walker County residents (about 85% of the population) have received type I oral poliomyelitis vaccine in a mass immunization program, utilizing vaccine from the Communicable Disease Center epidemic reserve. Plans are currently being formulated to give vaccine in the counties contiguous to Walker County where poliomyelitis cases have occurred.

Paliamyelitis (Cumulated Weekly) 1st Through 32nd Week

	1963	1962	1961	1960	1959
Paralytic	128	275	274	777	1,892
Total	153	364	413	1,074	2,924
Paliamyelitis (Six Week Totals) 27th Through 32nd Week					
Paralytic	60	114	119	431	1,198
Total	73	157	174	605	1,898

Table 1. CASES OF SPECIFIED NOTIFIABLE DISEASES: UNITED STATES
(Cumulative totals include revised and delayed reports through previous week)

Disease	32nd Week			Cumulative First 32 weeks		
	Ended August 10, 1963	Ended August 11, 1962	Median 1958 - 1962	1963	1962	Median 1958 - 1962
Aseptic meningitis	57	122	---	911	1,074	---
Brucellosis	9	7	11	225	257	481
Diphtheria	1	6	9	153	240	378
Encephalitis, infectious	19	46	52	958	987	999
Hepatitis, infectious and serum	802	734	645	27,597	36,160	22,896
Measles	1,528	1,660	1,746	354,109	437,864	390,448
Meningococcal infections	22	34	30	1,636	1,418	1,534
Poliomyelitis, total	12	30	127	153	364	1,076
Paralytic	11	21	97	128	275	776
Nonparalytic	-	9	24	15	63	225
Unspecified	1	-	6	10	26	75
Streptococcal sore throat and Scarlet fever	3,363	3,426	---	236,002	217,229	---
Tetanus	7	9	---	152	155	---
Tularemia	5	8	---	175	185	---
Typhoid fever	15	18	29	263	333	450
Typhus fever, tick-borne, (Rocky Mountain spotted)	6	17	---	110	152	---
Rabies in Animals	48	60	60	2,423	2,602	2,449

Table 2. NOTIFIABLE DISEASES OF LOW FREQUENCY

	Cum.		Cum.
Anthrax:	2	Psittacosis: Ill. - 1	39
Botulism:	5	Rabies in Man:	-
Malaria: N.Y. - 1, Md. - 1, Hawaii - 1, S. C. - 1	57	Smallpox:	-
Plague:	-	Typhus, murine:	14

Figures for current week exclude missing report from Alaska

SUMMARY OF REPORTED CASES OF INFECTIOUS SYPHILIS

JULY 1963 AND JULY 1962

CASES OF PRIMARY AND SECONDARY SYPHILIS: By Reporting Area July 1963 and July 1962 - Provisional Data

Reporting Area	July		Cumulative Jan - July	
	1963	1962	1963	1962
NEW ENGLAND	45	37	260	311
Maine	0	0	3	6
New Hampshire	1	1	5	7
Vermont	1	0	2	0
Massachusetts	37	25	156	215
Rhode Island	0	3	8	15
Connecticut	6	8	86	68
MIDDLE ATLANTIC	462	480	3,416	3,278
Upstate New York	70	53	367	324
New York City	226	288	1,892	1,849
Pa. (Excl. Phila.)	13	17	87	92
Philadelphia	53	57	425	412
New Jersey	100	65	645	601
EAST NORTH CENTRAL	143	158	1,117	976
Ohio	31	21	216	169
Indiana	6	11	36	51
Downstate Illinois	8	12	71	83
Chicago	73	81	524	479
Michigan	21	29	235	165
Wisconsin	4	4	35	29
WEST NORTH CENTRAL	29	32	276	214
Minnesota	6	9	46	35
Iowa	0	1	18	12
Missouri	17	18	127	109
North Dakota	0	0	4	2
South Dakota	1	0	18	18
Nebraska	2	1	29	6
Kansas	3	3	34	32
SOUTH ATLANTIC	568	437	3,531	3,259
Delaware	7	4	27	22
Maryland	67	41	333	310
District of Columbia	44	61	395	430
Virginia	25	31	176	231
West Virginia	2	6	26	27
North Carolina	92	58	531	370
South Carolina	95	50	419	446
Georgia	103	78	603	547
Florida	133	108	1,021	876
EAST SOUTH CENTRAL	139	112	903	663
Kentucky	19	14	85	77
Tennessee	33	40	234	187
Alabama	64	48	406	351
Mississippi	23	10	178	48
WEST SOUTH CENTRAL	247	291	1,663	1,702
Arkansas	13	31	131	137
Louisiana	48	121	348	736
Oklahoma	11	6	104	61
Texas	175	133	1,080	768
MOUNTAIN	31	17	274	203
Montana	2	1	5	1
Idaho	3	0	4	1
Wyoming	1	0	6	0
Colorado	1	1	23	41
New Mexico	9	5	70	34
Arizona	10	8	116	84
Utah	1	0	12	2
Nevada	4	2	38	40
PACIFIC	226	120	1,324	941
Washington	15	6	74	19
Oregon	4	3	32	28
California	204	110	1,201	881
Alaska	0	1	3	4
Hawaii	3	0	14	9
U. S. TOTAL	1,890	1,684	12,764	11,547
TERRITORIES	44	35	452	245
Puerto Rico	44	30	441	230
Virgin Islands	0	5	11	15

Note: Cumulative Totals include revised and delayed reports through previous months.

INFECTIOUS SYPHILIS – One thousand eight hundred and ninet cases of primary and secondary infectious syphilis were reported for the month of July 1963. This represents an increase of 12% as compared to July 1962, when 1,684 cases were reported. This rise is consistent with the upward trend of reported cases of infectious syphilis, a trend which has been evident in the United States for the past six years. The national per annum increases during the past two years have been about 10-12%.

With the exception of the New England and the West-South Central States, all geographic regions reported more cases during the period January-July 1963, than for the similar period of 1962. The decrease in reports of infectious syphilis in the West-South Central States is accounted for primarily by Louisiana where the 348 cases reported during January-July 1963, is less than one-half the 736 cases reported for the same period during 1962.

EPIDEMIOLOGICAL REPORT

Staphylococcal Food Poisoning – Ohio

On April 18, 1963, two men, both of whom had eaten ham sandwiches and coffee at a small restaurant in an Ohio town, suddenly developed marked pain and vomiting 1½ hours following lunch. On April 19, a third man developed similar symptoms 4½ hours following ingestion of a ham sandwich at this same restaurant. Two others, also known to have eaten ham sandwiches April 19, did not develop illness. In addition, the five employees of this restaurant also consumed the ham on April 18 without be coming ill.

The ham served at these meals had been deboned and excess fat trimmed off before baking on the morning of April 18. Allowed to cool, it was sliced before serving. An undetermined number of sandwiches were prepared from this ham, but no additional cases were uncovered on further investigation.

Samples of this ham as well as throat and stool cultures from all five employees were obtained for bacteriologic studies. Coagulase positive staphylococci, type 83A, were recovered both from the ham and the throat of one waitress. An untypable staphylococcus was cultured from another waitress' throat. Untypable staphylococci were recovered from the stools of four of the five employees.

The three men who became ill recovered rapidly, but no specimens were obtained for culture from them.

(Reported by Martha Laffey, RN, Findlay Health Commissioner; Division of Communicable Diseases and Division of Laboratories, Ohio Department of Health; and an EIS Officer from the Communicable Disease Center).

SUMMARY OF DEATHS AMONG PERSONS 65 YEARS AND OVER IN 108 U. S. CITIES

The weekly average number of deaths among persons 65 years and over in 108 cities for the four-week period ending August 10 was 6,070 as compared with an expected weekly average of 5,956.

	WEEK ENDING				4 Week Total	Weekly Average
	7/20	7/27	8/3	8/10		
Observed	6,197	6,141	6,217	5,725	24,280	6,070
Expected	5,982	5,959	5,944	5,937	23,822	5,956
Excess	215	182	273	-212	458	114

DEATHS AT AGE 65 AND OVER IN 108 U.S. CITIES
Average number per week by four-week periods



(See table, page 267)

Table 3 CASES OF SPECIFIED NOTIFIABLE DISEASES UNITED STATES
FOR WEEKS ENDED

AUGUST 10, 1963 AND AUGUST 11, 1962

Area	Poliomyelitis, total cases				Poliomyelitis, paralytic				Poliomyelitis, nonparalytic		Aseptic Meningitis	
	32nd week		Cumulative First 32 weeks		32nd week		Cumulative First 32 weeks		32nd week		32nd week	
	1963	1962	1963	1962	1963	1962	1963	1962	1963	1962	1963	1962
UNITED STATES	12	30	153	364	11	21	128	275	-	9	57	122
NEW ENGLAND	-	-	1	5	-	-	1	5	-	-	2	2
Maine	-	-	-	-	-	-	-	-	-	-	-	-
New Hampshire	-	-	-	-	-	-	-	-	-	-	-	-
Vermont	-	-	-	-	-	-	-	-	-	-	-	-
Massachusetts	-	-	1	4	-	-	1	4	-	-	1	1
Rhode Island	-	-	-	-	-	-	-	-	-	-	1	1
Connecticut	-	-	-	1	-	-	-	1	-	-	-	-
MIDDLE ATLANTIC	-	2	37	41	-	1	27	27	-	1	3	2
New York	-	2	7	37	-	1	5	23	-	1	2	2
New Jersey	-	-	1	3	-	-	1	3	-	-	-	-
Pennsylvania	-	-	29	1	-	-	21	1	-	-	1	-
EAST NORTH CENTRAL	3	5	23	32	2	5	15	21	-	-	12	9
Ohio	-	-	7	8	-	-	3	7	-	-	-	5
Indiana	-	1	2	5	-	1	1	4	-	-	4	1
Illinois	2	1	9	12	2	1	8	5	-	-	5	-
Michigan	-	3	2	5	-	3	2	5	-	-	3	3
Wisconsin	1	-	3	2	-	-	1	-	-	-	-	-
WEST NORTH CENTRAL	-	-	5	15	-	-	5	8	-	-	1	7
Minnesota	-	-	3	3	-	-	3	3	-	-	1	6
Iowa	-	-	-	5	-	-	-	2	-	-	-	1
Missouri	-	-	1	5	-	-	1	2	-	-	-	-
North Dakota	-	-	-	2	-	-	-	1	-	-	-	-
South Dakota	-	-	-	-	-	-	-	-	-	-	-	-
Nebraska	-	-	1	-	-	-	1	-	-	-	-	-
Kansas	-	-	-	-	-	-	-	-	-	-	-	-
SOUTH ATLANTIC	3	4	21	23	3	3	18	18	-	1	10	12
Delaware	-	-	-	-	-	-	-	-	-	-	1	-
Maryland	-	-	-	-	-	-	-	-	-	-	-	-
District of Columbia	-	-	-	1	-	-	-	-	-	-	-	-
Virginia	-	2	2	5	-	2	1	5	-	-	2	4
West Virginia	-	-	1	3	-	-	1	3	-	-	-	-
North Carolina	-	-	3	2	-	-	3	2	-	-	-	-
South Carolina	2	-	5	1	2	-	4	1	-	-	-	1
Georgia	1	2	5	4	1	1	4	3	-	1	-	-
Florida	-	-	5	7	-	-	5	4	-	-	7	7
EAST SOUTH CENTRAL	5	1	29	17	5	1	27	12	-	-	5	3
Kentucky	-	-	-	9	-	-	-	6	-	-	3	2
Tennessee	-	1	4	4	-	1	4	2	-	-	-	-
Alabama	5	-	23	3	5	-	21	3	-	-	2	1
Mississippi	-	-	2	1	-	-	2	1	-	-	-	-
WEST SOUTH CENTRAL	1	17	21	185	1	10	21	144	-	7	11	42
Arkansas	1	1	3	3	1	1	3	3	-	-	2	2
Louisiana	-	-	14	10	-	-	14	8	-	-	-	-
Oklahoma	-	-	-	3	-	-	-	1	-	-	-	-
Texas	-	16	4	169	-	9	4	132	-	7	9	40
MOUNTAIN	-	-	-	10	-	-	-	8	-	-	1	3
Montana	-	-	-	3	-	-	-	2	-	-	-	-
Idaho	-	-	-	1	-	-	-	1	-	-	-	-
Wyoming	-	-	-	1	-	-	-	1	-	-	-	-
Colorado	-	-	-	1	-	-	-	-	-	-	1	2
New Mexico	-	-	-	-	-	-	-	-	-	-	-	-
Arizona	-	-	-	3	-	-	-	3	-	-	-	-
Utah	-	-	-	1	-	-	-	1	-	-	-	1
Nevada	-	-	-	-	-	-	-	-	-	-	-	-
PACIFIC	-	1	16	36	-	1	14	32	-	-	12	42
Washington	-	-	-	-	-	-	-	-	-	-	1	-
Oregon	-	-	2	5	-	-	1	5	-	-	-	-
California	-	1	14	31	-	1	13	27	-	-	11	42
Alaska	---	-	-	-	---	-	-	-	---	-	---	-
Hawaii	-	-	-	-	-	-	-	-	-	-	-	-
Puerto Rico	-	-	4	9	-	-	4	9	-	-	-	-

Table 3. CASES OF SPECIFIED NOTIFIABLE DISEASES: UNITED STATES
FOR WEEKS ENDED
AUGUST 10, 1963 AND AUGUST 11, 1962 - (Continued)

Area	Brucellosis		Diphtheria		Encephalitis, infectious		Hepatitis, infectious and serum				Measles	
							32nd week					
	32nd week	Cumulative 32 weeks	32nd week	Cumulative 32 weeks	32nd week		Under 20 yr.	20 & over	Total		32nd week	
	1963	1963	1963	1963	1963	1962	1963	1963	1963	1962	1963	1962
UNITED STATES	9	225	1	153	19	46	390	368	802	734	1,528	1,660
NEW ENGLAND	1	1	-	8	1	3	52	44	103	62	88	113
Maine	-	-	-	-	-	-	28	19	47	21	1	12
New Hampshire	-	-	-	-	-	-	10	10	20	13	-	1
Vermont	1	1	-	-	-	-	2	1	3	2	5	1
Massachusetts	-	-	-	6	1	-	8	9	19	15	34	55
Rhode Island	-	-	-	2	-	3	-	-	2	-	5	12
Connecticut	-	-	-	-	-	-	4	5	12	11	43	32
MIDDLE ATLANTIC	-	5	-	21	2	12	88	72	160	135	255	231
New York	-	3	-	13	2	12	50	39	89	70	166	169
New Jersey	-	-	-	3	-	-	6	16	22	24	51	52
Pennsylvania	-	2	-	5	-	-	32	17	49	41	38	10
EAST NORTH CENTRAL	1	19	1	23	2	4	74	56	138	124	409	354
Ohio	-	-	-	1	-	2	23	11	36	37	90	23
Indiana	-	4	-	5	-	-	3	9	13	7	22	7
Illinois	1	12	1	12	1	2	21	13	35	36	49	48
Michigan	-	2	-	3	1	-	26	22	48	40	171	155
Wisconsin	-	1	-	2	-	-	1	1	6	4	77	121
WEST NORTH CENTRAL	5	138	-	37	1	2	13	12	27	43	42	65
Minnesota	-	8	-	15	-	-	3	2	5	5	1	12
Iowa	4	100	-	1	-	-	1	7	8	5	17	18
Missouri	1	10	-	1	-	2	3	2	7	15	-	8
North Dakota	-	-	-	1	1	-	1	-	1	3	24	23
South Dakota	-	8	-	11	-	-	1	-	1	1	-	4
Nebraska	-	6	-	8	-	-	-	1	1	9	-	-
Kansas	-	6	-	-	-	-	4	-	4	5	NN	NN
SOUTH ATLANTIC	2	12	-	31	5	7	29	31	61	107	216	124
Delaware	-	-	-	-	-	-	-	-	-	-	1	1
Maryland	-	-	-	-	1	-	2	4	6	12	31	8
District of Columbia	-	-	-	-	-	-	-	1	1	5	-	1
Virginia	1	4	-	-	1	-	5	6	11	18	25	47
West Virginia	-	-	-	1	-	-	4	3	7	10	53	32
North Carolina	1	4	-	2	-	1	16	3	19	34	-	1
South Carolina	-	-	-	9	-	1	-	1	2	10	9	2
Georgia	-	3	-	10	-	-	1	-	1	4	11	-
Florida	-	1	-	9	3	5	1	13	14	14	86	32
EAST SOUTH CENTRAL	-	10	-	10	-	2	36	25	61	79	74	74
Kentucky	-	3	-	-	-	-	17	7	24	22	28	8
Tennessee	-	5	-	2	-	1	9	7	16	38	38	59
Alabama	-	2	-	8	-	-	7	7	14	6	1	3
Mississippi	-	-	-	-	-	1	3	4	7	13	7	4
WEST SOUTH CENTRAL	-	25	-	17	1	4	33	28	62	65	109	123
Arkansas	-	4	-	1	1	1	2	5	7	13	1	-
Louisiana	-	6	-	2	-	-	9	11	20	3	1	-
Oklahoma	-	5	-	6	-	-	2	2	4	1	8	-
Texas	-	10	-	8	-	3	20	10	31	48	99	123
MOUNTAIN	-	7	-	4	-	2	27	26	76	28	130	187
Montana	-	-	-	-	-	-	1	4	7	2	12	61
Idaho	-	-	-	-	-	-	-	-	7	2	11	15
Wyoming	-	1	-	-	-	-	-	-	-	-	-	-
Colorado	-	-	-	3	-	-	1	-	7	8	31	36
New Mexico	-	-	-	1	-	-	2	1	3	5	NN	NN
Arizona	-	3	-	-	-	1	-	-	8	7	45	49
Utah	-	3	-	-	-	1	23	21	44	3	29	26
Nevada	-	-	-	-	-	-	-	-	-	1	2	-
PACIFIC	-	8	-	2	7	10	38	74	114	91	205	389
Washington	-	-	-	-	-	-	5	10	17	21	15	28
Oregon	-	2	-	-	-	1	1	6	7	6	21	48
California	-	5	-	2	7	9	32	56	88	62	161	167
Alaska	---	-	---	-	---	-	---	---	---	2	---	34
Hawaii	-	1	-	-	-	-	-	2	2	-	8	112
Puerto Rico	-	-	-	10	-	-	10	3	13	23	7	63

Table 3 CASES OF SPECIFIED NOTIFIABLE DISEASES UNITED STATES FOR WEEKS ENDED AUGUST 10, 1963 AND AUGUST 11, 1962 - (Continued)

Area	Meningococcal Infections		Streptococcal Sore Throat & Scarlet Fever		Tetanus	Tickborne Typhus (Rcky Mt. Spotted)	Tularemia	Typhoid Fever		Rabies in Animals		
	32nd wk.	Cumulative 32 weeks	32nd week		32nd wk.	32nd wk.	32nd wk.	32nd wk.	Cumulative 32 weeks	32nd week		Cumulative 32 weeks
	1963	1963	1963	1962	1963	1963	1963	1963	1963	1963	1962	1963
UNITED STATES	22	1,636	3,363	3,426	7	6	5	15	263	48	60	2,423
NEW ENGLAND	2	103	247	101	1	-	-	1	7	-	-	25
Maine	-	17	63	4	-	-	-	-	-	-	-	1
New Hampshire	-	4	7	2	-	-	-	-	-	-	-	12
Vermont	-	3	2	1	-	-	-	-	1	-	-	11
Massachusetts	2	50	30	19	1	-	-	-	4	-	-	1
Rhode Island	-	9	19	9	-	-	-	-	-	-	-	-
Connecticut	-	20	126	66	-	-	-	1	2	-	-	-
MIDDLE ATLANTIC	5	228	83	83	1	-	-	1	57	1	-	75
New York	4	107	68	68	1	-	-	-	19	1	-	55
New Jersey	-	32	9	10	-	-	-	-	3	-	-	-
Pennsylvania	1	89	6	5	-	-	-	1	35	-	-	20
EAST NORTH CENTRAL	1	253	195	195	3	1	-	1	18	12	17	408
Ohio	-	73	15	10	-	1	-	1	8	8	10	239
Indiana	-	31	50	48	3	-	-	-	1	-	3	40
Illinois	-	43	46	44	-	-	-	-	6	3	-	60
Michigan	1	79	55	52	-	-	-	-	1	-	4	39
Wisconsin	-	27	29	41	-	-	-	-	2	1	-	30
WEST NORTH CENTRAL	2	100	124	82	1	-	1	1	17	11	9	609
Minnesota	1	21	9	5	1	-	-	-	3	3	2	148
Iowa	-	5	29	28	-	-	-	-	2	4	1	234
Missouri	1	33	-	4	-	-	1	1	10	-	1	106
North Dakota	-	5	79	44	-	-	-	-	-	-	-	23
South Dakota	-	5	6	1	-	-	-	-	1	3	4	69
Nebraska	-	21	-	-	-	-	-	-	1	-	-	17
Kansas	-	10	1	-	-	-	-	-	-	1	1	12
SOUTH ATLANTIC	6	309	421	362	1	4	-	1	41	5	9	345
Delaware	-	2	-	-	-	1	-	-	1	-	-	1
Maryland	-	47	9	6	-	1	-	1	7	-	-	-
Dist. of Columbia	-	6	-	-	-	-	-	-	-	-	-	-
Virginia	-	72	62	77	1	1	-	-	5	2	-	125
West Virginia	-	16	112	77	-	1	-	-	6	2	7	100
North Carolina	-	52	3	8	-	-	-	-	6	-	-	6
South Carolina	1	15	33	8	-	-	-	-	2	-	-	6
Georgia	-	23	2	87	-	-	-	-	2	1	-	45
Florida	5	76	200	99	-	-	-	-	12	-	2	62
EAST SOUTH CENTRAL	1	122	657	852	-	1	-	4	26	3	3	200
Kentucky	1	26	46	14	-	-	-	-	1	1	-	95
Tennessee	-	53	567	755	-	1	-	4	18	2	2	88
Alabama	-	21	7	17	-	-	-	-	4	-	1	17
Mississippi	-	22	37	66	-	-	-	-	3	-	-	-
WEST SOUTH CENTRAL	3	159	474	598	-	-	2	3	42	7	10	464
Arkansas	1	11	-	1	-	-	2	-	18	3	1	40
Louisiana	2	65	4	1	-	-	-	3	11	-	-	38
Oklahoma	-	29	19	-	-	-	-	-	3	-	1	36
Texas	-	54	451	596	-	-	-	-	10	4	8	350
MOUNTAIN	-	55	741	673	-	-	2	2	13	4	-	61
Montana	-	3	33	19	-	-	-	-	-	-	-	-
Idaho	-	4	27	99	-	-	-	-	-	-	-	-
Wyoming	-	4	7	6	-	-	2	1	1	-	-	-
Colorado	-	14	303	235	-	-	-	1	6	1	-	7
New Mexico	-	4	211	178	-	-	-	-	2	-	-	21
Arizona	-	9	41	99	-	-	-	-	4	3	-	33
Utah	-	14	119	37	-	-	-	-	-	-	-	-
Nevada	-	3	-	-	-	-	-	-	-	-	-	-
PACIFIC	2	307	421	480	-	-	-	1	42	5	12	236
Washington	1	25	51	60	-	-	-	-	1	-	-	-
Oregon	-	22	4	7	-	-	-	-	2	-	2	1
California	1	241	320	387	-	-	-	1	36	5	10	226
Alaska	---	11	---	11	---	---	---	---	1	---	-	9
Hawaii	-	8	46	15	-	-	-	-	2	-	-	-
Puerto Rico	-	5	17	1	1	-	-	-	11	-	-	10

Table 4 (C). TOTAL DEATHS UNDER 1 YEAR OF AGE IN REPORTING CITIES

(Tables 4(A), 4(B), 4(C), and 4(D) will be published in sequence covering a four-week period.)°

Area	For weeks ending			
	7/20	7/27	8/3	8/10
NEW ENGLAND:				
Boston, Mass.	133	124	183	144
Bridgeport, Conn.	31	14	24	23
Cambridge, Mass.	22	17	23	18
Fall River, Mass.	14	16	19	10
Hartford, Conn.	39	22	39	16
Lowell, Mass.	12	14	28	24
Lynn, Mass.	19	13	30	9
New Bedford, Mass.	17	16	26	17
New Haven, Conn.	32	25	30	26
Providence, R.I.	27	36	45	27
Somerville, Mass.	12	11	14	9
Springfield, Mass.	32	29	32	26
Waterbury, Conn.	13	24	20	11
Worcester, Mass.	22	31	46	26
MIDDLE ATLANTIC:				
Albany, N.Y.	21	19	28	18
Allentown, Pa.	15	21	17	15
Buffalo, N.Y.	72	77	89	51
Camden, N.J.	19	24	19	17
Elizabeth, N.J.	17	12	17	14
Erie, Pa.	25	26	19	19
Jersey City, N.J.	44	40	43	35
Newark, N.J.	59	35	42	41
New York City, N.Y.	859	865	992	762
Paterson, N.J.	23	15	26	20
Philadelphia, Pa.	316	304	301	345
Pittsburgh, Pa.	128	107	101	98
Reading, Pa.	27	29	31	38
Rochester, N.Y.	67	52	78	63*
Schenectady, N.Y.	18	15	17	17
Scranton, Pa.	19	28	17	18
Syracuse, N.Y.	32	42	32	30
Trenton, N.J.	19	38	16	25
Utica, N.Y.	25	22	13	10
Yonkers, N.Y.	19	15	19	16
EAST NORTH CENTRAL:				
Akron, Ohio	27	23	27	30
Canton, Ohio	20	16	12	22
Chicago, Ill.	368	339	351	343
Cincinnati, Ohio	94	76	92	96
Cleveland, Ohio	98	110	102	112
Columbus, Ohio	54	63	68	55
Dayton, Ohio	56	49	44	47
Detroit, Mich.	167	160	204	173
Evansville, Ind.	23	27	22	19
Flint, Mich.	14	24	28	22
Fort Wayne, Ind.	29	19	24	27
Gary, Ind.	9	8	20	17
Grand Rapids, Mich.	26	25	21	31
Indianapolis, Ind.	73	86	51	90
Madison, Wis.	13	13	21	21
Milwaukee, Wis.	64	50	63	74
Peoria, Ill.	16	15	9	15
Rockford, Ill.	12	16	26	19
South Bend, Ind.	34	24	23	14
Toledo, Ohio	43	53	62	51
Youngstown, Ohio	37	39	31	36
WEST NORTH CENTRAL:				
Des Moines, Iowa	32	31	29	31
Duluth, Minn.	21	7	21	13
Kansas City, Kans.	14	28	20	18
Kansas City, Mo.	100	80	66	78
Lincoln, Nebr.	16	19	13	20
Minneapolis, Minn.	72	83	84	66
Omaha, Nebr.	54	43	37	32
St. Louis, Mo.	126	124	115	119
St. Paul, Minn.	38	40	52	37
Wichita, Kans.	44	47	26	31
SOUTH ATLANTIC:				
Atlanta, Ga.	56	44	43	54
Baltimore, Md.	137	141	118	91
Charlotte, N.C.	19	20	13	17
Jacksonville, Fla.	24	28	22	37
Miami, Fla.	43	47	47	42
Norfolk, Va.	30	29	21	25
Richmond, Va.	49	36	52	49
Savannah, Ga.	5	12	12	9
St. Petersburg, Fla.	44	42	37	59
Tampa, Fla.	39	39	31	32
Washington, D.C.	77	95	79	76
Wilmington, Del.	25	28	10	22
EAST SOUTH CENTRAL:				
Birmingham, Ala.	33	42	35	35
Chattanooga, Tenn.	16	34	16	13
Knoxville, Tenn.	15	21	18	13
Louisville, Ky.	51	70	62	41
Memphis, Tenn.	46	49	66	53
Mobile, Ala.	24	14	13	24
Montgomery, Ala.	13	23	15	11
Nashville, Tenn.	46	59	45	43
WEST SOUTH CENTRAL:				
Austin, Tex.	29	24	18	19
Baton Rouge, La.	17	15	9	6
Corpus Christi, Tex.	11	12	17	12
Dallas, Tex.	59	76	60	44
El Paso, Tex.	19	14	19	12
Fort Worth, Tex.	45	42	35	27
Houston, Tex.	93	82	72	73
Little Rock, Ark.	37	34	38	25
New Orleans, La.	85	81	85	93
Oklahoma City, Okla.	35	49	47	42
San Antonio, Tex.	51	46	48	45
Shreveport, La.	25	24	37	31
Tulsa, Okla.	30	24	21	36
MOUNTAIN:				
Albuquerque, N. Mex.	17	22	7	12
Colorado Springs, Colo.	11	16	6	12
Denver, Colo.	64	75	54	57
Ogden, Utah	2	9	14	10
Phoenix, Ariz.	34	67	45	43
Pueblo, Colo.	10	5	10	11
Salt Lake City, Utah	41	28	34	35
Tucson, Ariz.	21	25	26	14
PACIFIC:				
Berkeley, Calif.	6	13	12	10
Fresno, Calif.	24	18	33	25
Glendale, Calif.	28	20	18	20
Honolulu, Hawaii	18	18	21	17
Long Beach, Calif.	39	44	34	27
Los Angeles, Calif.	309	293	278	271
Oakland, Calif.	59	51	40	55
Pasadena, Calif.	22	17	19	25*
Portland, Oreg.	65	76	65	65
Sacramento, Calif.	31	29	28	31*
San Diego, Calif.	56	43	39	49
San Francisco, Calif.	85	91	87	108
San Jose, Calif.	21	20	15	19
Seattle, Wash.	79	77	87	78*
Spokane, Wash.	34	32	29	26
Tacoma, Wash.	25	33	18	31
San Juan, P.R.	18	12	13	(---)

*Estimate - based on average percent of divisional total.
Totals for previous weeks include reported corrections.

NOTE: All deaths by place of occurrence.

°Current Week Mortality for 108 Selected Cities

4(A) Total Mortality, all ages	10,620
4(B) Pneumonia-Influenza Deaths, all ages	361
4(C) Total Deaths under 1 Year of Age	702
4(D) Total Deaths, Persons 65 years and over	5,725

UNIVERSITY OF FLORIDA
3 1262 08864 0312

UNIV OF FL LIB
DOCUMENTS DEPT
U.S. DEPOSITORY

INTERNATIONAL NOTES

Smallpox – Sweden

Sweden declared Stockholm free of smallpox as of August 5, 1963. The total number of cases reported in this outbreak was 26, with 4 deaths. (Previous reports appeared in MMWR, Vol. 12, pp. 174, 183, and 236.)

Smallpox – Poland

No new cases have been reported in Wroclaw (Breslau) or elsewhere in Poland, according to the information received from the World Health Organization as of August 12. (Previous reports appeared in MMWR, Vol. 12, pp. 244, 252, and 255.)

QUARANTINE MEASURES

Immunization Information for International Travel
1962 Edition
Public Health Service Publication No. 384

Section 6, Page 69 – Under Los Angeles, California, delete all information concerning the U. S. Public Health Service Outpatient Clinic.

ERRATUM

In the Hepatitis Morbidity Report published last week (Vol. 12, No. 31, page 254), it was incorrectly stated that Louisiana was among those States in which hepatitis rates were higher in 1963 than during the three previous years. The peak year for Louisiana occurred in 1961-1962.

In addition to the established procedures for reporting morbidity and mortality, the Communicable Disease Center welcomes accounts of interesting outbreaks or cases. Such accounts should be addressed to:

Editor, Morbidity and Mortality Weekly Report
Communicable Disease Center
Atlanta, Georgia 30333

U. S. DEPARTMENT OF
HEALTH, EDUCATION, AND WELFARE
PUBLIC HEALTH SERVICE
Communicable Disease Center
Atlanta, Georgia 30333

Official Business

POSTAGE AND FEES PAID
U. S. DEPARTMENT OF H. E. W.